


VECTORS IN 2 DIMENSIONS


➤ **Vector notation:** _____

➤ **Magnitude of a vector** - given by the Pythagorean theorem.

- notation: _____




➤ **Direction of a vector** - given by the inverse of the tangent ratio of the vertical and horizontal vector components.



➤ **Vector components** are also vectors.

• **Horizontal vector component:**





• **Vertical vector component:**



Example 1: Sketch a labeled diagram of a displacement vector $\vec{d} = 14 \text{ m [N}28^\circ\text{W]}$. Find the components of the vector and rewrite the vector using vector notation.

Example 2: Determine the magnitude and direction of each vector. Write the information about each vector in two different ways.






Practice:

1. Given the diagram of vector \vec{m} , determine the following:

- Vector notation of \vec{m} : _____
- Vector components of \vec{m} : _____ or _____
_____ or _____
- Magnitude of \vec{m} : _____
- Direction of \vec{m} : _____



2. Determine vector components of $\vec{b} = 13\text{cm}[L75^\circ D]$. Include a diagram in your solution.

3. Classify the given quantities as either a scalar or a vector.

Mass	
Volume	
Force	
Potential energy	
Work	
Kinetic energy	
Time	
Displacement	
Temperature	
Distance	
Height	

Momentum	
Speed	
Position	
Velocity	
Efficiency	
Power	
Acceleration	
Tension	
Impulse	
Weight	
Length	

Review Math for Physics

- Conversion of units using dimensional analysis

Example 1: Express 25 m/s in km/h

Example 2: Express density of 3.5 g/cm³ in kg/m³

- **Rounding:** carry as many digits throughout your calculations as you reasonably can. Round at the very end and determine the number of digits needed based on the rules for significant digits and operations with numbers.
- **Addition and subtraction:** The result of addition or subtraction is permitted to have the same number of decimal places as the number with the least number of decimal places.
- **Multiplication and division:** The result of multiplication or division is permitted to have the same number of decimal places as the number with the least number of significant digits.

- Scientific notation: use it for all final answers.

- Significant digits
 - Non-zero digits are also significant

 - Zeros between non-zero digits are significant

 - Zeros on the right of the decimal point when there is at least one non-zero digit to the left of them are significant

 - Zeros on the right of the decimal point when there are no non-zero digits to the left of them are not significant

 - Zeros at the end of an integer (=non-decimal number) may or may not be significant.